

The effect of the Galactic gas distribution on the expected Cosmic Rays spectrum

Mercedes Mollá, Manuel Aguilar, Juan Alcaraz, Javier Berdugo, Jorge Casaus, Carlos Díaz, Elisa Lanciotti, Carlos Mañá, Jesús Marín, Gustavo Martínez, Carmen Palomares, Eusebio Sánchez, Ignacio Sevilla, and Ana Sofía Torrentó

CIEMAT, Avda. Complutense 22, 28040 Madrid (Spain)

1 Introduction

Cosmic ray (CR) nuclei are accelerated particles which move randomly through the interstellar medium (ISM), where they suffer scattering, reacceleration and energy loss processes before reaching Earth. Spallation processes also take place forming secondary nuclei by fragmentation of heavier ones. Due to the impossibility of observing directly their original direction, the determination of possible sources where these particles originated requires the use of codes to simulate the propagation of CR within the Galaxy. This consists of a spiral disk with a thickness of $2h \sim 200$ pc, where CR are created, and a halo with a height H , where they diffuse.

From the existing data, mostly $^{10}\text{Be}/^9\text{Be}$ and B/C , it is deduced that CR go through a mean density of $\sim 0.3 \text{ cm}^{-3}$ ([4,5]). Since the total density of the ISM is considered as 1 cm^{-3} , ($\langle n_{\text{HI}} \rangle \sim 0.5 \text{ cm}^{-3}$; [1,2,7,11]), a large effective halo and/or a local bubble or cavity of low density are required. This result is very dependent on the actual disk ISM density, largely uncertain. [6] have very recently obtained a new map of the diffuse gas distribution (Fig. 1), from which $\langle n_{\text{HI}} \rangle \sim 0.2 \text{ cm}^{-3}$, a factor of 2 lower than previous estimates.




Fig. 1. The radial distribution of the diffuse gas atomic density as obtained from [6] compared with other author's distributions as labeled in the figure

2 Computed Models and Results

We now use the GALPROP code, developed by [8,9], to compute how CR propagates. The model includes a realistic diffuse gas radial distribution ([3]). A halo height $H = 1 - 5$ kpc is obtained, depending of the model. Results for $^{10}\text{Be}/^9\text{Be}$ and B/C are shown with dashed lines in Fig. 2 a) and b), respectively, for a diffusion model with $H = 1$ kpc.

We now change the old HI distribution with the new one. The dotted lines in Fig.2 represent the same model using this new distribution. It is evident that it is not yet valid. The solid lines correspond to a new good enough model, with a halo height $H = 0.5$ kpc. This implies that CR diffuse mostly in the disk closer to the sources than it was deduced before. This result agrees with [10] who obtain distances as short as 500 pc between the source and the Sun, from the ACE radionuclide measurements.

References

1. Dickey, J. M. & Lockman, F.J. 1990, A.R.A.A., **28**, 215
2. Ferrière, K. 1998, ApJ., **497**, 759
3. Gordon, M.A. & Burton, W.B. 1976, ApJ., **208**, 346
4. Jones, F. C., Lukasiak, A., Ptuskin, V., & Webber, W. 2001, ApJ, **547**, 264
5. Maurin, D., Donato, F., Taillet, R. & Salati, P. 2001, ApJ., **555**, 585
6. Nakanishi, H. & Sofue, Y. 2004, PASJ, **55**, 191
7. Olling, R. P. & Merrifield, M. R. 2001, MNRAS, **326**, 164
8. Strong & Moskalenko 1998, ApJ, **509**, 212
9. Strong & Moskalenko 2003, 28 IRC Conf., 1921
10. Yanasak, N.E., Wiedenbeck, M. E., Mewaldt, R.A. et al. 2001, ApJ., **563**, 768
11. Wolfire, M. G., McKee, C. F., Hollenbach, D., et al. 2003, ApJ, **587**, 27




Fig. 2. The spectra of a) $^{10}\text{Be}/^9\text{Be}$, and b) of B/C. Models are represented by lines as labeled